

# EXEMPLAR SOLUTIONS MATHS

Class  
**11**



Powered by



## Chapter 3- Trigonometric Functions

### EXERCISE

#### SHORT ANSWER TYPE

1. Prove that

$$\frac{\tan A + \sec A - 1}{\tan A - \sec A + 1} = \frac{1 + \sin A}{\cos A}$$

**Solution:**

According to the question,

$$\begin{aligned} \text{LHS} &= \frac{\tan A + \sec A - 1}{\tan A - \sec A + 1} \\ &= \frac{\frac{\sin A}{\cos A} + \frac{1}{\cos A} - 1}{\frac{\sin A}{\cos A} - \frac{1}{\cos A} + 1} \\ &= \frac{\frac{\sin A + 1 - \cos A}{\cos A}}{\frac{\sin A - 1 + \cos A}{\cos A}} \\ &= \frac{\sin A + 1 - \cos A}{\sin A - 1 + \cos A} \end{aligned}$$

Using the identity,

$\sin^2 A + \cos^2 A = 1$ , we get,

$\sin A + (1 - \cos A)$ .

$$\begin{aligned} \therefore \text{LHS} &= \frac{\sin A + (1 - \cos A)}{\sin A - (1 - \cos A)} \times \frac{\sin A + (1 - \cos A)}{\sin A + (1 - \cos A)} \\ &= \frac{\{\sin A + (1 - \cos A)\}^2}{\sin^2 A - (1 - \cos A)^2} \\ &= \frac{\sin^2 A - (1 - \cos A)^2}{\sin^2 A + (1 - \cos A)^2 + 2 \sin A (1 - \cos A)} \\ &= \frac{\sin^2 A - (1 - \cos A)^2}{(\sin^2 A + \cos^2 A) + 1 - 2 \cos A + 2 \sin A (1 - \cos A)} \\ &= \frac{\sin^2 A - \{1 + \cos^2 A - 2 \cos A\}}{(1) + 1 - 2 \cos A + 2 \sin A (1 - \cos A)} \\ &= \frac{(\sin^2 A - 1) - \cos^2 A + 2 \cos A}{2(1 - \cos A) + 2 \sin A (1 - \cos A)} \\ &= \frac{(-\cos^2 A) - \cos^2 A + 2 \cos A}{2(1 + \sin A)(1 - \cos A)} \\ &= \frac{-2 \cos^2 A + 2 \cos A}{2(1 + \sin A)(1 - \cos A)} \\ &= \frac{2 \cos A (1 - \cos A)}{(1 + \sin A) \cos A} = \text{RHS} \end{aligned}$$

Hence, L.H.S = R.H.S



2. If  $[2\sin\alpha / (1+\cos\alpha+\sin\alpha)] = y$ , then prove that  $[(1-\cos\alpha+\sin\alpha) / (1+\sin\alpha)]$  is also equal to y.

[Hint : Express  $\frac{1-\cos\alpha+\sin\alpha}{1+\sin\alpha} = \frac{1-\cos\alpha+\sin\alpha}{1+\sin\alpha} \cdot \frac{1+\cos\alpha+\sin\alpha}{1+\cos\alpha+\sin\alpha}$ ]

**Solution:**

According to the question,

$$y = 2\sin\alpha / (1+\cos\alpha+\sin\alpha)$$

Multiplying numerator and denominator by  $(1-\cos\alpha+\sin\alpha)$ ,

We get,

$$\begin{aligned} \Rightarrow y &= \frac{2\sin\alpha}{1+\cos\alpha+\sin\alpha} \times \frac{1-\cos\alpha+\sin\alpha}{1-\cos\alpha+\sin\alpha} \\ &= \frac{2\sin\alpha}{(1+\sin\alpha)+\cos\alpha} \times \frac{(1+\sin\alpha)-\cos\alpha}{(1+\sin\alpha)-\cos\alpha} \end{aligned}$$

Using  $(a+b)(a-b) = a^2 - b^2$ , we get:

$$\begin{aligned} &= \frac{2\sin\alpha \{(1+\sin\alpha) - \cos\alpha\}}{(1+\sin\alpha)^2 - \cos^2\alpha} \\ &= \frac{2\sin\alpha (1+\sin\alpha) - 2\sin\alpha \cos\alpha}{1 + \sin^2\alpha + 2\sin\alpha - \cos^2\alpha} \end{aligned}$$

Since,  $1 - \cos^2\alpha = \sin^2\alpha$

$$\begin{aligned} \therefore y &= \frac{2\sin\alpha (1+\sin\alpha - \cos\alpha)}{\sin^2\alpha + 2\sin\alpha + \sin^2\alpha} \\ &= \frac{2\sin\alpha (1+\sin\alpha - \cos\alpha)}{2\sin\alpha (1+\sin\alpha)} \\ \Rightarrow y &= \frac{(1+\sin\alpha - \cos\alpha)}{(1+\sin\alpha)} \\ \Rightarrow y &= \frac{(1-\cos\alpha+\sin\alpha)}{(1+\sin\alpha)} \end{aligned}$$

Hence Proved

3. If  $m \sin \theta = n \sin (\theta + 2\alpha)$ , then prove that

$$\tan (\theta + \alpha) \cot \alpha = (m + n)/(m - n)$$

[Hints: Express  $\sin(\theta + 2\alpha) / \sin\theta = m/n$  and apply componendo and dividend]

**Solution:**

According to the question,

$$m \sin \theta = n \sin (\theta + 2\alpha)$$

To prove:

$$\tan (\theta + \alpha) \cot \alpha = (m + n)/(m - n)$$

Proof:

$$m \sin \theta = n \sin (\theta + 2\alpha)$$

$$\Rightarrow \sin(\theta + 2\alpha) / \sin \theta = m/n$$

Applying componendo-dividendo rule, we have,

$$\frac{\sin(\theta+2\alpha)+\sin \theta}{\sin(\theta+2\alpha)-\sin \theta} = \frac{m+n}{m-n}$$

By transformation formula of T-ratios,

We know that,

$$\sin A + \sin B = 2 \sin ((A+B)/2) \cos ((A - B)/2)$$

And,

$$\sin A - \sin B = 2 \cos ((A+B)/2) \sin ((A - B)/2)$$

On applying the formula, we get,

$$\frac{2 \sin \left( \frac{2\theta + 2\alpha}{2} \right) \cos \left( \frac{\theta + 2\alpha - \theta}{2} \right)}{2 \cos \left( \frac{2\theta + 2\alpha}{2} \right) \sin \left( \frac{\theta + 2\alpha - \theta}{2} \right)} = \frac{m+n}{m-n}$$

$$\frac{\sin(\theta+\alpha) \cos(\alpha)}{\cos(\theta+\alpha) \sin(\alpha)} = \frac{m+n}{m-n}$$

$$\{\because \tan x = (\sin x)/(\cos x)\}$$

$$\Rightarrow \tan(\theta + \alpha) \cot \alpha = \frac{m+n}{m-n}$$

$$\text{Therefore, } \tan (\theta + \alpha) \cot \alpha = (m + n)/(m - n)$$

Hence Proved

$$\cos(\alpha + \beta) = \frac{4}{5} \text{ and } \sin(\alpha - \beta) = \frac{5}{13},$$

4. If  
2α

where α lie between 0 and π/4, find value of tan

[Hint: Express tan 2α as tan (α + β + α - β)]

**Solution:**

According to the question,

$$\cos(\alpha + \beta) = 4/5 \dots(i)$$

We know that,

$$\sin x = \sqrt{1 - \cos^2 x}$$

Therefore,

$$\sin (\alpha + \beta) = \sqrt{1 - \cos^2(\alpha + \beta)}$$

$$\Rightarrow \sin (\alpha + \beta) = \sqrt{1 - (4/5)^2} = 3/5 \dots(ii)$$

Also,

$$\sin(\alpha - \beta) = 5/13 \text{ \{given\} } \dots(iii)$$

we know that,

$$\cos x = \sqrt{1 - \sin^2 x}$$

Therefore,

$$\cos (\alpha - \beta) = \sqrt{1 - \sin^2(\alpha - \beta)}$$

$$\Rightarrow \cos (\alpha - \beta) = \sqrt{1 - (5/13)^2} = 12/13 \dots(iv)$$

Therefore,

$$\tan 2\alpha = \tan (\alpha + \beta + \alpha - \beta)$$

We know that,

$$\tan(x+y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

$$\tan(\alpha+\beta) + \tan(\alpha-\beta)$$

$$\therefore \tan 2\alpha = \frac{1 - \tan(\alpha+\beta) \tan(\alpha-\beta)}{\frac{\sin(\alpha+\beta)}{\cos(\alpha+\beta)} + \frac{\sin(\alpha-\beta)}{\cos(\alpha-\beta)}}$$

$$\Rightarrow \tan 2\alpha = \frac{1 - \frac{\sin(\alpha+\beta)}{\cos(\alpha+\beta)} \times \frac{\sin(\alpha-\beta)}{\cos(\alpha-\beta)}}{\frac{\sin(\alpha+\beta)}{\cos(\alpha+\beta)} + \frac{\sin(\alpha-\beta)}{\cos(\alpha-\beta)}}$$

From equation i, ii, iii and iv we have,

$$\Rightarrow \tan 2\alpha = \frac{\frac{3}{5} + \frac{5}{13}}{1 - \frac{3}{5} \times \frac{5}{13}}$$

$$= \frac{\frac{3}{4} + \frac{5}{12}}{1 - \frac{3}{4} \times \frac{5}{12}}$$

$$= \frac{\frac{9+5}{12}}{1 - \frac{15}{48}}$$

$$\Rightarrow \tan 2\alpha = \frac{14}{12 \left( \frac{33}{48} \right)}$$

$$= \frac{56}{33}$$

Hence,  $\tan 2\alpha = 56/33$

5. If  $\tan x = b/a$  then find the value of

$$\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}$$

**Solution:**

According to the question,

$$\tan x = b/a$$

Let,

**NAGEEN GROUP**  
tamso ma jyotirgamaya

$$\begin{aligned}
 y &= \sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}} \\
 \therefore y &= \sqrt{\frac{a(1+\frac{b}{a})}{a(1-\frac{b}{a})}} + \sqrt{\frac{a(1-\frac{b}{a})}{a(1+\frac{b}{a})}} \\
 &= \sqrt{\frac{(1+\tan x)}{(1-\tan x)}} + \sqrt{\frac{(1-\tan x)}{(1+\tan x)}} \\
 &= \frac{\sqrt{1+\tan x}}{\sqrt{1-\tan x}} + \frac{\sqrt{1-\tan x}}{\sqrt{1+\tan x}} \\
 &= \frac{(\sqrt{1+\tan x})^2 + (\sqrt{1-\tan x})^2}{(\sqrt{1-\tan x})(\sqrt{1+\tan x})} \\
 &= \frac{1+\tan x + 1-\tan x}{\sqrt{1-\tan^2 x}} = \frac{2}{\sqrt{1-\tan^2 x}} \\
 \therefore y &= \frac{\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}}{2} = \frac{2}{\sqrt{1-\tan^2 x}} \\
 &= \frac{\sqrt{1-\frac{\sin^2 \theta}{\cos^2 \theta}}}{2} \\
 &= \frac{\sqrt{\cos^2 \theta - \sin^2 \theta}}{2 \cos \theta} \\
 \therefore \cos^2 \theta - \sin^2 \theta &= \cos 2\theta \\
 &= \frac{2 \cos \theta}{\sqrt{\cos 2\theta}}
 \end{aligned}$$

**6. Prove that  $\cos \theta \cos \theta/2 - \cos 3\theta \cos 9\theta/2 = \sin 7\theta \sin 4\theta$**

**[Hint: Express L.H.S. =  $\frac{1}{2} [2\cos \theta \cos \theta/2 - 2\cos 3\theta \cos 9\theta / 2]$**

**Solution:**

Using transformation formula, we get,

$$2 \cos A \cos B = \cos(A + B) + \cos(A - B)$$

$$-2 \sin A \sin B = \cos(A + B) - \cos(A - B)$$

Multiplying and dividing the expression by 2.

**INAGEEN GROUP**  
tamso ma jyotirgamaya

$$\therefore \text{LHS} = \frac{1}{2} \left( 2 \cos \theta \cos \frac{\theta}{2} - 2 \cos 3\theta \cos \frac{9\theta}{2} \right)$$

Applying transformation formula, we get,

$$\text{LHS} = \frac{1}{2} \left( \cos \left( \theta + \frac{\theta}{2} \right) + \cos \left( \theta - \frac{\theta}{2} \right) - \left\{ \cos \left( 3\theta + \frac{9\theta}{2} \right) + \cos \left( 3\theta - \frac{9\theta}{2} \right) \right\} \right)$$

$$\Rightarrow \text{LHS} = \frac{1}{2} \left( \cos \frac{3\theta}{2} + \cos \frac{\theta}{2} - \cos \left( \frac{15\theta}{2} \right) - \cos \left( -\frac{3\theta}{2} \right) \right)$$

$$\Rightarrow \text{LHS} = \frac{1}{2} \left( \cos \frac{3\theta}{2} + \cos \frac{\theta}{2} - \cos \frac{15\theta}{2} - \cos \frac{3\theta}{2} \right) \{ \because \cos(-x) = \cos x \}$$

$$\Rightarrow \text{LHS} = \frac{1}{2} \left( \cos \frac{\theta}{2} - \cos \frac{15\theta}{2} \right)$$

$$\Rightarrow \text{LHS} = \frac{1}{2} \left( 2 \sin \left( \frac{\frac{\theta}{2} + \frac{15\theta}{2}}{2} \right) \sin \left( \frac{\frac{15\theta}{2} - \frac{\theta}{2}}{2} \right) \right)$$

$$\Rightarrow \text{LHS} = \frac{1}{2} \left( 2 \sin \left( \frac{8\theta}{2} \right) \sin \left( \frac{7\theta}{2} \right) \right)$$

$$\therefore \text{LHS} = \sin 4\theta \sin \left( \frac{7\theta}{2} \right) = \text{RHS}$$

Hence,

$$\cos \theta \cos \frac{\theta}{2} - \cos 3\theta \cos \frac{9\theta}{2} = \sin 4\theta \sin \left( \frac{7\theta}{2} \right)$$

**7. If  $a \cos \theta + b \sin \theta = m$  and  $a \sin \theta - b \cos \theta = n$ , then show that  $a^2 + b^2 = m^2 + n^2$ .**

**Solution:**

According to the question,

$$a \cos \theta + b \sin \theta = m \dots (i)$$

$$a \sin \theta - b \cos \theta = n \dots (ii)$$

Squaring and adding equation 1 and 2, we get,

$$(a \cos \theta + b \sin \theta)^2 + (a \sin \theta - b \cos \theta)^2 = m^2 + n^2$$

$$\Rightarrow a^2 \cos^2 \theta + b^2 \sin^2 \theta + 2ab \sin \theta \cos \theta + a^2 \sin^2 \theta + b^2 \cos^2 \theta - 2ab \sin \theta \cos \theta = m^2 + n^2$$

$$\Rightarrow a^2 \cos^2 \theta + b^2 \sin^2 \theta + a^2 \sin^2 \theta + b^2 \cos^2 \theta = m^2 + n^2$$

$$\Rightarrow a^2 (\sin^2 \theta + \cos^2 \theta) + b^2 (\sin^2 \theta + \cos^2 \theta) = m^2 + n^2$$

Using,  $\sin^2 \theta + \cos^2 \theta = 1$ ,

We get,

$$\Rightarrow a^2 + b^2 = m^2 + n^2$$

**8. Find the value of  $\tan 22^\circ 30'$ .**

$$\tan \frac{\theta}{2} = \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}} = \frac{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}} = \frac{\sin \theta}{1 + \cos \theta}$$

[Hint: Let  $\theta = 45^\circ$ , use

**Solution:**

Let,  $\theta = 45^\circ$

As we need to find:  $\tan 22^\circ 30' = \tan (\theta/2)$

We know that,

$$\sin \theta = \cos \theta = 1/\sqrt{2} \text{ (for } \theta = 45^\circ \text{)}$$

Since,



$$\tan \frac{\theta}{2} = \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}}$$

Multiplying  $2\cos \frac{\theta}{2}$  in numerator and denominator, we get,

$$\Rightarrow \tan \frac{\theta}{2} = \frac{2 \cos \frac{\theta}{2} \sin \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}}$$

By applying formula of T-ratios of multiple angles-

$$\sin 2x = 2 \sin x \cos x$$

$$\cos 2x = 2\cos^2 x - 1 \text{ or } 1 + \cos 2x = 2\cos^2 x$$

$$\therefore \tan \frac{\theta}{2} = \frac{\sin \theta}{1 + \cos \theta}$$

$$\Rightarrow \tan 22^\circ 30' = \frac{\sin 45^\circ}{1 + \cos 45^\circ}$$

$$= \frac{\frac{1}{\sqrt{2}}}{1 + \frac{1}{\sqrt{2}}}$$

$$= \frac{1}{\sqrt{2} + 1}$$

$$\text{Therefore, } \tan 22^\circ 30' = \sqrt{2} - 1$$

**9. Prove that  $\sin 4A = 4\sin A \cos^3 A - 4\cos A \sin^3 A$ .**

**Solution:**

$$\sin 4A = \sin (2A + 2A)$$

We know that,

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\text{Therefore, } \sin 4A = \sin 2A \cos 2A + \cos 2A \sin 2A$$

$$\Rightarrow \sin 4A = 2 \sin 2A \cos 2A$$

From T-ratios of multiple angle,

We get,

$$\sin 2A = 2 \sin A \cos A \text{ and } \cos 2A = \cos^2 A - \sin^2 A$$

$$\Rightarrow \sin 4A = 2(2 \sin A \cos A)(\cos^2 A - \sin^2 A)$$

$$\Rightarrow \sin 4A = 4 \sin A \cos^3 A - 4 \cos A \sin^3 A$$

$$\text{Hence, } \sin 4A = 4 \sin A \cos^3 A - 4 \cos A \sin^3 A$$

**10. If  $\tan \theta + \sin \theta = m$  and  $\tan \theta - \sin \theta = n$ , then prove that  $m^2 - n^2 = 4 \sin \theta \tan \theta$**

**[Hint:  $m + n = 2 \tan \theta$ ,  $m - n = 2 \sin \theta$ , then use  $m^2 - n^2 = (m + n)(m - n)$ ]**

**Solution:**

According to the question,

$$\tan \theta + \sin \theta = m \dots (i)$$

$$\tan \theta - \sin \theta = n \dots (ii)$$

Adding equation i and ii,

$$2 \tan \theta = m + n \dots (iii)$$

Subtracting equation ii from i,

We get,

$$2\sin \theta = m - n \dots(\text{iv})$$

Multiplying equations (iii) and (iv),

$$2\sin \theta (2\tan \theta) = (m + n)(m - n)$$

$$\Rightarrow 4 \sin \theta \tan \theta = m^2 - n^2$$

Hence,

$$m^2 - n^2 = 4 \sin \theta \tan \theta$$

**11. If  $\tan (A + B) = p$ ,  $\tan (A - B) = q$ , then show that  $\tan 2A = (p + q) / (1 - pq)$ .**

**[Hint: Use  $2A = (A + B) + (A - B)$ ]**

**Solution:**

We know that,

$$\tan 2A = \tan (A + B + A - B)$$

And also,

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

$$\therefore \tan 2A = \frac{\tan(A+B) + \tan(A-B)}{1 - \tan(A+B) \tan(A-B)}$$

Substituting the values given in question,

$$\Rightarrow \tan 2A = \frac{p+q}{1-pq}$$

$$\text{Hence, } \tan 2A = \frac{p+q}{1-pq}$$

**12. If  $\cos \alpha + \cos \beta = 0 = \sin \alpha + \sin \beta$ , then prove that  $\cos 2\alpha + \cos 2\beta = -2\cos (\alpha + \beta)$ .**

**[Hint:  $(\cos \alpha + \cos \beta)^2 - (\sin \alpha + \sin \beta)^2 = 0$ ]**

**Solution:**

According to the question,

$$\cos \alpha + \cos \beta = 0 = \sin \alpha + \sin \beta \dots(\text{i})$$

Since, LHS =  $\cos 2\alpha + \cos 2\beta$

We know that,

$$\cos 2x = \cos^2 x - \sin^2 x$$

Therefore,

$$\text{LHS} = \cos^2 \alpha - \sin^2 \alpha + (\cos^2 \beta - \sin^2 \beta)$$

$$\Rightarrow \text{LHS} = \cos^2 \alpha + \cos^2 \beta - (\sin^2 \alpha + \sin^2 \beta)$$

Also, since,

$$a^2 + b^2 = (a+b)^2 - 2ab$$

$$\Rightarrow \text{LHS} = (\cos \alpha + \cos \beta)^2 - 2\cos \alpha \cos \beta - (\sin \alpha + \sin \beta)^2 + 2\sin \alpha \sin \beta$$

From equation (i),

$$\Rightarrow \text{LHS} = 0 - 2\cos \alpha \cos \beta - 0 + 2\sin \alpha \sin \beta$$

$$\Rightarrow \text{LHS} = -2(\cos \alpha \cos \beta - \sin \alpha \sin \beta)$$

$$\because \cos (\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\text{Therefore, LHS} = -2 \cos (\alpha + \beta) = \text{RHS}$$

$$\text{Hence, } \cos 2\alpha + \cos 2\beta = -2\cos (\alpha + \beta)$$

**13.**

If  $\frac{\sin(x+y)}{\sin(x-y)} = \frac{a+b}{a-b}$ , then show that  $\frac{\tan x}{\tan y} = \frac{a}{b}$ .

[Hint: Use componendo and Dividendo]

**Solution:**

According to the question,

$$\frac{\sin(x+y)}{\sin(x-y)} = \frac{a+b}{a-b}$$

$$\frac{\sin(x+y)}{\sin(x-y)} = \frac{a+b}{a-b}$$

Since,  $\sin(A+B) = \sin A \cos B + \cos A \sin B$

$$\frac{\sin(x+y)}{\sin(x-y)} = \frac{a+b}{a-b}$$

$$\therefore \frac{\sin(x+y)}{\sin(x-y)} = \frac{a+b}{a-b}$$

$$\frac{\sin x \cos y + \cos x \sin y}{\sin x \cos y - \cos x \sin y} = \frac{a+b}{a-b}$$

$$\Rightarrow \frac{\sin x \cos y + \cos x \sin y}{\sin x \cos y - \cos x \sin y} = \frac{a+b}{a-b}$$

Applying componendo-dividendo rule,

We get,

$$\frac{(\sin x \cos y + \cos x \sin y) + (\sin x \cos y - \cos x \sin y)}{(\sin x \cos y + \cos x \sin y) - (\sin x \cos y - \cos x \sin y)} = \frac{(a+b) + (a-b)}{(a+b) - (a-b)}$$

$$\Rightarrow \frac{2 \sin x \cos y}{2 \cos x \sin y} = \frac{2a}{2b}$$

$$\Rightarrow \frac{2 \sin x \cos y}{2 \cos x \sin y} = \frac{2a}{2b}$$

$$\Rightarrow \left( \frac{\sin x}{\cos x} \right) \left( \frac{\cos y}{\sin y} \right) = \frac{a}{b}$$

$$\Rightarrow \left( \frac{\sin x}{\cos x} \right) \left( \frac{\cos y}{\sin y} \right) = \frac{a}{b}$$

$$\Rightarrow \left( \frac{\sin x}{\cos x} \right) \left( \frac{\cos y}{\sin y} \right) = \frac{a}{b}$$

Since,  $\tan A = (\sin A)/(\cos A)$

$$\Rightarrow \tan x \left( \frac{1}{\tan y} \right) = \frac{a}{b}$$

$$\Rightarrow \tan x \left( \frac{1}{\tan y} \right) = \frac{a}{b}$$

$$\Rightarrow \frac{\tan x}{\tan y} = \frac{a}{b}$$

$$\Rightarrow \frac{\tan x}{\tan y} = \frac{a}{b}$$

14.

$$\tan \theta = \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha},$$

If  $\tan \theta = \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha}$ , then show that  $\sin \alpha + \cos \alpha = \sqrt{2} \cos \theta$ .

[Hint: Express  $\tan \theta = \tan(\alpha - \pi/2)$   $\theta = \alpha - \pi/4$ ]

**Solution:**

We know that,

$$\tan \theta = \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha}$$

$$\tan \theta = \frac{\cos \alpha \left( \frac{\sin \alpha}{\cos \alpha} - 1 \right)}{\cos \alpha \left( \frac{\sin \alpha}{\cos \alpha} + 1 \right)}$$

Since,  $\tan A = (\sin A)/(\cos A)$

$$\Rightarrow \tan \theta = (\tan \alpha - 1) / (\tan \alpha + 1)$$

Since,  $\tan \pi/4 = 1$

$$\tan \theta = \frac{(\tan \alpha - \tan \frac{\pi}{4})}{(1 + \tan \frac{\pi}{4} \cdot \tan \alpha)}$$

We know that,

$$\tan(x-y) = (\tan x - \tan y) / (1 + \tan x \cdot \tan y)$$

$$\text{Therefore, } \tan \theta = \tan (\alpha - \pi/4)$$

$$\Rightarrow \theta = \alpha - \pi/4$$

$$\Rightarrow \alpha = \theta + \pi/4 \dots (i)$$

To prove,

$$\sin \alpha + \cos \alpha = \sqrt{2} \cos \theta$$

$$\because \text{LHS} = \sin \alpha + \cos \alpha$$

From equation (i)

$$\Rightarrow \text{LHS} = \sin(\theta + \pi/4) + \cos(\theta + \pi/4)$$

$$\because \sin(x+y) = \sin x \cos y + \cos x \sin y$$

$$\text{And, } \cos(x+y) = \cos x \cos y - \sin x \sin y$$

$$\text{Therefore, LHS} = \sin \theta \cos(\pi/4) + \sin(\pi/4) \cos \theta + \cos \theta \cos(\pi/4) - \sin(\pi/4) \sin \theta$$

$$\because \sin(\pi/4) = \cos(\pi/4) = 1/\sqrt{2}$$

$$\Rightarrow \text{LHS} = \sin \theta (1/\sqrt{2}) + (1/\sqrt{2}) \cos \theta + \cos \theta (1/\sqrt{2}) - \sin \theta (1/\sqrt{2})$$

$$\Rightarrow \text{LHS} = 2 \cos \theta (1/\sqrt{2})$$

$$\Rightarrow \text{LHS} = \sqrt{2} \cos \theta = \text{RHS}$$

$$\text{Therefore, } \sin \alpha + \cos \alpha = \sqrt{2} \cos \theta$$

**15. If  $\sin \theta + \cos \theta = 1$ , then find the general value of  $\theta$ .**

**Solution:**

According to the question,

$$\sin \theta + \cos \theta = 1$$

$$\text{As, } \sin \theta + \cos \theta = 1$$

$$\Rightarrow \sqrt{2} \left( \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta \right) = 1$$

We know that,

$$\sin(\pi/4) = \cos(\pi/4) = 1/\sqrt{2}$$

$$\Rightarrow \sqrt{2} \left( \sin \theta \cos \frac{\pi}{4} + \sin \frac{\pi}{4} \cos \theta \right) = 1$$

We know that,

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\Rightarrow \sin \left( \frac{\pi}{4} + \theta \right) = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \sin \left( \frac{\pi}{4} + \theta \right) = \sin \frac{\pi}{4}$$

Since we know,

$$\text{If } \sin \theta = \sin \alpha \Rightarrow \theta = n\pi + (-1)^n \alpha$$

We get,

$$\theta + \pi/4 = n\pi + (-1)^n (\pi/4)$$

$$\Rightarrow \theta = n\pi + (\pi/4)((-1)^n - 1)$$

**16. Find the most general value of  $\theta$  satisfying the equation  $\tan \theta = -1$  and  $\cos \theta = 1/\sqrt{2}$**

**Solution:**

According to the question,

We have,

$$\tan \theta = -1$$

$$\text{And } \cos \theta = 1/\sqrt{2}.$$

$$\Rightarrow \theta = -\pi/4$$

So, we know that,

$\theta$  lies in IV quadrant.

$$\theta = 2\pi - \pi/4 = 7\pi/4$$

So, general solution is  $\theta = 7\pi/4 + 2n\pi, n \in \mathbb{Z}$

**17. If  $\cot \theta + \tan \theta = 2 \operatorname{cosec} \theta$ , then find the general value of  $\theta$ .**

**Solution:**

According to the question,

$$\Rightarrow \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{\cos \theta} = 2 \operatorname{cosec} \theta$$

Since,

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\Rightarrow \frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta \cos \theta} = 2 \operatorname{cosec} \theta$$

$$\Rightarrow 1 = 2 \operatorname{cosec} \theta \sin \theta \cos \theta$$

We know that,

$$\sin \theta \operatorname{cosec} \theta = 1$$

$$\Rightarrow 1 = 2 \cos \theta$$

$$\Rightarrow \cos \theta = 1/2 = \cos(\pi/3)$$

Hence,

The solution of  $\cos x = \cos \alpha$  can be given by,

$$x = 2m\pi \pm \alpha \quad \forall m \in \mathbb{Z}$$

$$\Rightarrow \theta = 2n\pi \pm \pi/3, n \in \mathbb{Z}$$

**18. If  $2\sin^2 \theta = 3\cos \theta$ , where  $0 \leq \theta \leq 2\pi$ , then find the value of  $\theta$ .**

**Solution:**

According to the question,

$$2\sin^2 \theta = 3\cos \theta$$

We know that,

$$\sin^2 \theta = 1 - \cos^2 \theta$$

Given that,

$$2\sin^2 \theta = 3\cos \theta$$

$$2 - 2\cos^2 \theta = 3\cos \theta$$

$$2\cos^2 \theta + 3\cos \theta - 2 = 0$$

$$(\cos \theta + 2)(2\cos \theta - 1) = 0$$

Therefore,

$$\cos \theta = 1/2 = \cos \pi/3$$

$$\theta = \pi/3 \text{ or } 2\pi - \pi/3$$

$$\theta = \pi/3, 5\pi/3$$

Therefore,  $2(1 - \cos^2 \theta) = 3\cos \theta$

$$\Rightarrow 2 - 2\cos^2 \theta = 3\cos \theta$$

$$\Rightarrow 2\cos^2 \theta + 3\cos \theta - 2 = 0$$

$$\Rightarrow 2\cos^2 \theta + 4\cos \theta - \cos \theta - 2 = 0$$



$$\Rightarrow 2\cos \theta (\cos \theta + 2) + 1 (\cos \theta + 2) = 0$$

$$\Rightarrow (2\cos \theta + 1)(\cos \theta + 2) = 0$$

Since,  $\cos \theta \in [-1, 1]$ , for any value  $\theta$ .

So,  $\cos \theta \neq -2$

Therefore,

$$2\cos \theta - 1 = 0$$

$$\Rightarrow \cos \theta = \frac{1}{2}$$

$$= \pi/3 \text{ or } 2\pi - \pi/3$$

$$\theta = \pi/3, 5\pi/3$$

**19. If  $\sec x \cos 5x + 1 = 0$ , where  $0 < x \leq \pi/2$ , then find the value of  $x$ .**

**Solution:**

According to the question,

$$\sec x \cos 5x = -1$$

$$\Rightarrow \cos 5x = -1/\sec x$$

We know that,

$$\sec x = 1/\cos x$$

$$\Rightarrow \cos 5x + \cos x = 0$$

By transformation formula of T-ratios,

We know that,

$$\cos A + \cos B = 2\cos \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right)$$

$$\Rightarrow 2\cos \left( \frac{5x+x}{2} \right) \cos \left( \frac{5x-x}{2} \right) = 0$$

$$\Rightarrow 2\cos 3x \cos 2x = 0$$

$$\Rightarrow \cos 3x = 0 \text{ or } \cos 2x = 0$$

$$\because 0 < x \leq \pi/2$$

$$\text{Therefore, } 0 < 2x \leq \pi \text{ or } 0 < 3x \leq 3\pi/2$$

$$\text{Therefore, } 2x = \pi/2$$

$$\Rightarrow x = \pi/4$$

$$3x = \pi/2$$

$$\Rightarrow x = \pi/6$$

$$\text{Or } 3x = 3\pi/2$$

$$\Rightarrow x = \pi/2$$

$$\text{Hence, } x = \pi/6, \pi/4, \pi/2.$$

**20. If  $\sin (\theta + \alpha) = a$  and  $\sin (\theta + \beta) = b$ , then prove that  $\cos 2(\alpha - \beta) - 4ab \cos (\alpha - \beta) = 1 - 2a^2 - 2b^2$**

**Solution:**

According to the question,

$$\sin (\theta + \alpha) = a \text{ and } \sin (\theta + \beta) = b$$

$$\text{LHS} = \cos 2(\alpha - \beta) - 4ab \cos (\alpha - \beta)$$

$$\text{Using } \cos 2x = 2\cos^2 x - 1,$$

Let us solve,

$$\Rightarrow \text{LHS} = 2\cos^2(\alpha - \beta) - 1 - 4ab \cos(\alpha - \beta)$$

$$\Rightarrow \text{LHS} = 2\cos(\alpha - \beta) \{\cos(\alpha - \beta) - 2ab\} - 1$$

Since,

$$\cos(\alpha - \beta) = \cos\{(\theta + \alpha) - (\theta + \beta)\}$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\Rightarrow \cos(\alpha - \beta) = \cos(\theta + \alpha)\cos(\theta + \beta) + \sin(\theta + \alpha)\sin(\theta + \beta)$$

Since,

$$\sin(\theta + \alpha) = a$$

$$\Rightarrow \cos(\theta + \alpha) = \sqrt{1 - \sin^2(\theta + \alpha)} = \sqrt{1 - a^2}$$

Similarly,

$$\cos(\theta + \beta) = \sqrt{1 - b^2}$$

Therefore,

$$\cos(\alpha - \beta) = \sqrt{1 - a^2}\sqrt{1 - b^2} + ab$$

Therefore,

$$\text{LHS} = 2\{ab + \sqrt{1 - a^2}(1 - b^2)\} \{ab + \sqrt{1 - a^2}(1 - b^2) - 2ab\} - 1$$

$$\Rightarrow \text{LHS} = 2\{\sqrt{1 - a^2}(1 - b^2) + ab\} \{\sqrt{1 - a^2}(1 - b^2) - ab\} - 1$$

$$\text{Using } (x + y)(x - y) = x^2 - y^2$$

$$\Rightarrow \text{LHS} = 2\{(1 - a^2)(1 - b^2) - a^2b^2\} - 1$$

$$\Rightarrow \text{LHS} = 2\{1 - a^2 - b^2 + a^2b^2\} - 1$$

$$\Rightarrow \text{LHS} = 2 - 2a^2 - 2b^2 - 1$$

$$\Rightarrow \text{LHS} = 1 - 2a^2 - 2b^2 = \text{RHS}$$

Therefore,

We get,

$$\cos 2(\alpha - \beta) - 4ab \cos(\alpha - \beta) = 1 - 2a^2 - 2b^2$$

**21. If  $\cos(\theta + \phi) = m \cos(\theta - \phi)$ , then prove that  $\tan \theta = ((1 - m)/(1 + m)) \cot \phi$**

**[Hint: Express  $\cos(\theta + \phi)/\cos(\theta - \phi) = m/1$  and apply Componendo and Dividendo]**

**Solution:**

According to the question,

**NAGEEN GROUP**  
tamso ma jyotirgamaya

$$\begin{aligned}\cos(\theta + \phi) &= m \cos(\theta - \phi) \\ \therefore \cos(\theta + \phi) &= m \cos(\theta - \phi) \\ \frac{\cos(\theta - \phi)}{\cos(\theta + \phi)} &= \frac{1}{m}\end{aligned}$$

Applying componendo – dividend, we get,

$$\begin{aligned}\frac{\cos(\theta - \phi) + \cos(\theta + \phi)}{\cos(\theta - \phi) - \cos(\theta + \phi)} &= \frac{1+m}{1-m}\end{aligned}$$

From transformation formula, we know that,

$$\cos(A+B) + \cos(A-B) = 2\cos A \cos B$$

$$\cos(A-B) - \cos(A+B) = 2\sin A \sin B$$

$$\begin{aligned}\frac{2\cos\theta\cos\phi}{2\sin\theta\sin\phi} &= \frac{1+m}{1-m}\end{aligned}$$

Since,  $(\cos\theta)/(\sin\theta) = \cot\theta$

$$\Rightarrow \cot\theta \cot\phi = \frac{1+m}{1-m}$$

$$\Rightarrow \left(\frac{1-m}{1+m}\right) \cot\phi = \frac{1}{\cot\theta}$$

$$\Rightarrow \tan\theta = \left(\frac{1-m}{1+m}\right) \cot\phi$$

**22. Find the value of the expression**

$$3\left[\sin^4\left(\frac{3\pi}{2} - \alpha\right) + \sin^4(3\pi + \alpha)\right] - 2\left[\sin^6\left(\frac{\pi}{2} + \alpha\right) + \sin^6(5\pi - \alpha)\right]$$

**Solution:**

According to the question,

$$\text{Let, } y = 3[\sin^4(3\pi/2 - \alpha) + \sin^4(3\pi + \alpha)] - 2[\sin^6(\pi/2 + \alpha) + \sin^6(5\pi - \alpha)]$$

We know that,

$$\sin(3\pi/2 - \alpha) = -\cos\alpha$$

$$\sin(3\pi + \alpha) = -\sin\alpha$$

$$\sin(\pi/2 + \alpha) = \cos\alpha$$

$$\sin(5\pi - \alpha) = \sin\alpha$$

Therefore,

$$y = 3[(-\cos\alpha)^4 + (-\sin\alpha)^4] - 2[\cos^6\alpha + \sin^6\alpha]$$

$$\Rightarrow y = 3[\cos^4\alpha + \sin^4\alpha] - 2[\sin^6\alpha + \cos^6\alpha]$$

$$\Rightarrow y = 3[(\sin^2\alpha + \cos^2\alpha)^2 - 2\sin^2\alpha \cos^2\alpha] - 2[(\sin^2\alpha)^3 + (\cos^2\alpha)^3]$$

Since, we know that,

$$\sin^2\alpha + \cos^2\alpha = 1$$

Also, we know that,

$$a^3 + b^3 = (a+b)(a^2 - ab + b^2)$$

$$\Rightarrow y = 3[1 - 2\sin^2\alpha \cos^2\alpha] - 2[(\sin^2\alpha + \cos^2\alpha)(\cos^4\alpha + \sin^4\alpha - \sin^2\alpha \cos^2\alpha)]$$

$$\Rightarrow y = 3[1 - 2\sin^2\alpha \cos^2\alpha] - 2[\cos^4\alpha + \sin^4\alpha - \sin^2\alpha \cos^2\alpha]$$

$$\Rightarrow y = 3[1 - 2\sin^2\alpha \cos^2\alpha] - 2[(\sin^2\alpha + \cos^2\alpha)^2 - 2\sin^2\alpha \cos^2\alpha - \sin^2\alpha \cos^2\alpha]$$

$$\Rightarrow y = 3[1 - 2\sin^2\alpha \cos^2\alpha] - 2[1 - 3\sin^2\alpha \cos^2\alpha]$$

$$\Rightarrow y = 3 - 6\sin^2\alpha \cos^2\alpha - 2 + 6\sin^2\alpha \cos^2\alpha$$

$$\Rightarrow y = 1$$

**23. If  $a \cos 2\theta + b \sin 2\theta = c$  has  $\alpha$  and  $\beta$  as its roots, then prove that  $\tan \alpha + \tan \beta = 2b/(a + c)$   
[Hint: Use the identities  $\cos 2\theta = ((1 - \tan^2 \theta)/(1 + \tan^2 \theta))$  and  $\sin 2\theta = 2\tan \theta/(1 + \tan^2 \theta)$ ]**

**Solution:**

According to the question,

$$a \cos 2\theta + b \sin 2\theta = c$$

$\alpha$  and  $\beta$  are the roots of the equation.

Using the formula of multiple angles,

We know that,

$$\cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \text{ and } \sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta}$$

$$\therefore a \left( \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right) + b \left( \frac{2 \tan \theta}{1 + \tan^2 \theta} \right) - c = 0$$

$$\Rightarrow a(1 - \tan^2 \theta) + 2b \tan \theta - c(1 + \tan^2 \theta) = 0$$

$$\Rightarrow (-c - a)\tan^2 \theta + 2b \tan \theta - c + a = 0 \dots (i)$$

We know that,

The sum of roots of a quadratic equation,  $ax^2 + bx + c = 0$  is given by  $(-b/a)$

Therefore,

$$\tan \alpha + \tan \beta = -2b/(-c + a) = 2b/(c + a)$$

$$\text{Hence, } \tan \alpha + \tan \beta = 2b/(c + a)$$

**24. If  $x = \sec \phi - \tan \phi$  and  $y = \operatorname{cosec} \phi + \cot \phi$ , then show that  $xy + x - y + 1 = 0$ .**

**[Hint: Find  $xy + 1$  and then show  $\tan x - y = -(xy + 1)$ ]**

**Solution:**

According to the question,

$$x = \sec \phi - \tan \phi \text{ and } y = \operatorname{cosec} \phi + \cot \phi$$

Given that, LHS =  $xy + x - y + 1$

$$= (\sec \phi - \tan \phi)(\operatorname{cosec} \phi + \cot \phi) + (\sec \phi - \tan \phi) - (\operatorname{cosec} \phi + \cot \phi) + 1$$

$$= \sec \phi \operatorname{cosec} \phi + \cot \phi \sec \phi - \tan \phi \cot \phi - \tan \phi \operatorname{cosec} \phi$$

$$+ \sec \phi - \tan \phi - (\operatorname{cosec} \phi + \cot \phi) + 1$$

$$= \frac{1}{\sin \phi \cos \phi} + \frac{1}{\sin \phi} - 1 - \sec \phi + \sec \phi - \tan \phi - \left( \frac{1}{\sin \phi} + \frac{\cos \phi}{\sin \phi} \right) + 1$$

$$= \frac{1}{\sin \phi \cos \phi} + \frac{1}{\sin \phi} - \tan \phi - \left( \frac{1}{\sin \phi} + \frac{\cos \phi}{\sin \phi} \right)$$

$$= \frac{1}{\sin \phi \cos \phi} - \frac{\sin \phi}{\cos \phi} - \frac{\cos \phi}{\sin \phi}$$

$$= \frac{1}{\sin \phi \cos \phi} - \left( \frac{\cos \phi}{\sin \phi} + \frac{\sin \phi}{\cos \phi} \right)$$

$$= \frac{1}{\sin \phi \cos \phi} - \left( \frac{\cos^2 \phi + \sin^2 \phi}{\sin \phi \cos \phi} \right)$$

$$\text{Since, } \sin^2 \theta + \cos^2 \theta = 1$$

$$= \frac{1}{\sin \phi \cos \phi} - \left( \frac{1}{\sin \phi \cos \phi} \right) = 0$$

$$\text{Thus, LHS} = xy + x - y + 1 = 0$$

**25. If  $\theta$  lies in the first quadrant and  $\cos \theta = 8/17$ , then find the value of  $\cos(30^\circ + \theta) + \cos(45^\circ - \theta) + \cos(120^\circ - \theta)$**

**Solution:**

According to the question,

$$\cos \theta = 8/17$$

$$\sin \theta = \pm\sqrt{1 - \cos^2 \theta}$$

Since,  $\theta$  lies in first quadrant, only positive sign can be considered.

$$\Rightarrow \sin \theta = \sqrt{1 - 64/289} = 15/17$$

$$\text{Let, } y = \cos(30^\circ + \theta) + \cos(45^\circ - \theta) + \cos(120^\circ - \theta)$$

We know that,

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

Therefore,

$$y = \cos 30^\circ \cos \theta - \sin 30^\circ \sin \theta + \cos 45^\circ \cos \theta + \sin 45^\circ \sin \theta + \cos 120^\circ \cos \theta + \sin 120^\circ \sin \theta$$

Substituting values of  $\cos 30^\circ$ ,  $\sin 30^\circ$ ,  $\cos 120^\circ$ ,  $\sin 120^\circ$  and  $\cos 45^\circ$

$$\Rightarrow y = \frac{\sqrt{3}}{2} \cdot \frac{8}{17} - \frac{1}{2} \cdot \frac{15}{17} + \frac{1}{\sqrt{2}} \cdot \frac{8}{17} + \frac{1}{\sqrt{2}} \cdot \frac{15}{17} + \left(-\frac{1}{2}\right) \left(\frac{8}{17}\right) + \left(\frac{\sqrt{3}}{2}\right) \left(\frac{15}{17}\right)$$

$$= \frac{8\sqrt{3}}{34} - \frac{15}{34} + \frac{8 + 15}{17\sqrt{2}} - \frac{8}{34} + \frac{15\sqrt{3}}{34}$$

$$= \frac{23\sqrt{3}}{34} + \frac{23}{17\sqrt{2}} - \frac{23}{34}$$

$$\Rightarrow y = \frac{23}{34} (\sqrt{3} + \sqrt{2} - 1)$$

**26. Find the value of the expression  $\cos^4(\pi/8) + \cos^4(3\pi/8) + \cos^4(5\pi/8) + \cos^4(7\pi/8)$ .**

**[Hint: Simplify the expression to**

$$2 \left( \cos^4 \frac{\pi}{8} + \cos^4 \frac{3\pi}{8} \right) = 2 \left[ \left( \cos^2 \frac{\pi}{8} + \cos^2 \frac{3\pi}{8} \right)^2 - 2 \cos^2 \frac{\pi}{8} \cos^2 \frac{3\pi}{8} \right]$$

**Solution:**

According to the question,

$$\text{Let } y = \cos^4(\pi/8) + \cos^4(3\pi/8) + \cos^4(5\pi/8) + \cos^4(7\pi/8).$$

$$\Rightarrow y = \cos^4(\pi/8) + \cos^4(3\pi/8) + \cos^4(\pi - 3\pi/8) + \cos^4(\pi - \pi/8).$$

Since we know that,  $\cos(\pi - x) = -\cos x$ , we get,

**NAGEEN GROUP**  
tamso ma jyotirgamaya



$$\begin{aligned}
&= \cos^4 \frac{\pi}{8} + \cos^4 \frac{3\pi}{8} + \cos^4 \left( \frac{3\pi}{8} \right) + \cos^4 \left( \frac{\pi}{8} \right) \\
&= 2 \left( \cos^4 \frac{\pi}{8} + \cos^4 \frac{3\pi}{8} \right) \\
&= 2 \left( \cos^4 \frac{\pi}{8} + \cos^4 \left( \frac{\pi}{2} - \frac{\pi}{8} \right) \right) \\
&= 2 \left( \cos^4 \frac{\pi}{8} + \sin^4 \frac{\pi}{8} \right) \\
&= 2 \left[ \left( \cos^2 \frac{\pi}{8} + \sin^2 \frac{\pi}{8} \right)^2 - 2 \cos^2 \frac{\pi}{8} \cdot \sin^2 \frac{\pi}{8} \right] \\
&= 2 \left[ 1 - 2 \cos^2 \frac{\pi}{8} \cdot \sin^2 \frac{\pi}{8} \right] \\
&= 2 - \left( 2 \cos \frac{\pi}{8} \cdot \sin \frac{\pi}{8} \right)^2 \\
&= 2 - \left( \sin \frac{2\pi}{8} \right)^2 \\
&= 2 - (1/\sqrt{2})^2 \\
&= 2 - 1/2 \\
&= 3/2
\end{aligned}$$

**27. Find the general solution of the equation**

$$5\cos^2\theta + 7\sin^2\theta - 6 = 0$$

**Solution:**

According to the question,

$$5\cos^2\theta + 7\sin^2\theta - 6 = 0$$

We know that,

$$\sin^2\theta = 1 - \cos^2\theta$$

$$\text{Therefore, } 5\cos^2\theta + 7(1 - \cos^2\theta) - 6 = 0$$

$$\Rightarrow 5\cos^2\theta + 7 - 7\cos^2\theta - 6 = 0$$

$$\Rightarrow -2\cos^2\theta + 1 = 0$$

$$\Rightarrow \cos^2\theta = 1/2$$

$$\text{Therefore, } \cos \theta = \pm 1/\sqrt{2}$$

$$\text{Therefore, } \cos \theta = \cos \pi/4 \text{ or } \cos \theta = \cos 3\pi/4$$

Since, solution of  $\cos x = \cos \alpha$  is given by

$$x = 2m\pi \pm \alpha \quad \forall m \in \mathbb{Z}$$

$$\theta = n\pi \pm \pi/4, n \in \mathbb{Z}$$

**28. Find the general solution of the equation  $\sin x - 3\sin 2x + \sin 3x = \cos x - 3\cos 2x + \cos 3x$**

**Solution:**

According to the question,

$$\sin x - 3\sin 2x + \sin 3x = \cos x - 3\cos 2x + \cos 3x$$

Grouping  $\sin x$  and  $\sin 3x$  in LHS and,  $\cos x$  and  $\cos 3x$  in RHS,

We get,

$$\sin x + \sin 3x - 3\sin 2x = \cos x + \cos 3x - 3\cos 2x$$

Applying transformation formula,

$$\cos A + \cos B = 2\cos((A+B)/2) \cos((A-B)/2)$$

$$\begin{aligned} \sin A + \sin B &= 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \\ \Rightarrow 2 \sin \left( \frac{3x+x}{2} \right) \cos \left( \frac{3x-x}{2} \right) - 3 \sin 2x &= 2 \cos \left( \frac{3x+x}{2} \right) \cos \left( \frac{3x-x}{2} \right) - 3 \cos 2x \\ \Rightarrow 2 \sin 2x \cos x - 3 \sin 2x &= 2 \cos 2x \cos x - 3 \cos 2x \\ \Rightarrow 2 \sin 2x \cos x - 3 \sin 2x - 2 \cos 2x \cos x + 3 \cos 2x &= 0 \\ \Rightarrow 2 \cos x (\sin 2x - \cos 2x) - 3(\sin 2x - \cos 2x) &= 0 \\ \Rightarrow (\sin 2x - \cos 2x)(2 \cos x - 3) &= 0 \\ \Rightarrow \cos x = 3/2 \text{ or } \sin 2x = \cos 2x \\ \text{As } \cos x \in [-1, 1] \\ \text{Hence, no value of } x \text{ exists for which } \cos x = 3/2 \\ \text{Therefore, } \sin 2x = \cos 2x \\ \Rightarrow \tan 2x = 1 = \tan \pi/4 \\ \text{We know solution of } \tan x = \tan \alpha \text{ is given by,} \\ x = n\pi \pm \alpha, n \in \mathbb{Z} \\ \text{Therefore, } 2x = n\pi \pm (\pi/4) \\ \Rightarrow x = n\pi/2 \pm (\pi/8), n \in \mathbb{Z} \end{aligned}$$

**29. Find the general solution of the equation  $(\sqrt{3}-1) \cos \theta + (\sqrt{3}+1) \sin \theta = 2$**   
**[Hint: Put  $\sqrt{3}-1 = r \sin \alpha$ ,  $\sqrt{3}+1 = r \cos \alpha$  which gives  $\tan \alpha = \tan((\pi/4) - (\pi/6))$   $\alpha = \pi/12$ ]**

**Solution:**

$$\begin{aligned} \text{Let, } r \sin \alpha &= \sqrt{3}-1 \text{ and } r \cos \alpha = \sqrt{3}+1 \\ \text{Therefore, } r &= \sqrt{(\sqrt{3}-1)^2 + (\sqrt{3}+1)^2} = \sqrt{8} = 2\sqrt{2} \\ \text{And, } \tan \alpha &= (\sqrt{3}-1) / (\sqrt{3}+1) \\ \text{Therefore, } r(\sin \alpha \cos \theta + \cos \alpha \sin \theta) &= 2 \\ \Rightarrow r \sin (\theta + \alpha) &= 2 \\ \Rightarrow \sin (\theta + \alpha) &= 1/\sqrt{2} \\ \Rightarrow \sin (\theta + \alpha) &= \sin (\pi/4) \\ \Rightarrow \theta + \alpha &= n\pi + (-1)^n (\pi/4), n \in \mathbb{Z} \\ \Rightarrow \theta &= n\pi + (-1)^n (\pi/4) - (\pi/12), n \in \mathbb{Z} \end{aligned}$$

### OBJECTIVE TYPE QUESTIONS

**30. If  $\sin \theta + \operatorname{cosec} \theta = 2$ , then  $\sin^2 \theta + \operatorname{cosec}^2 \theta$  is equal to**

- A. 1
- B. 4
- C. 2

**D. None of these**

**Solution:**

C. 2

Explanation:

According to the question,

$$\sin \theta + \operatorname{cosec} \theta = 2$$

Squaring LHS and RHS,

We get,

$$\Rightarrow (\sin \theta + \operatorname{cosec} \theta)^2 = 2^2$$

$$\Rightarrow \sin^2 \theta + \operatorname{cosec}^2 \theta + 2 \sin \theta \operatorname{cosec} \theta = 4$$

$$\Rightarrow \sin^2\theta + \operatorname{cosec}^2\theta + 2 \sin \theta (1/\sin)\theta = 4$$

$$\Rightarrow \sin^2\theta + \operatorname{cosec}^2\theta + 2 = 4$$

$$\Rightarrow \sin^2\theta + \operatorname{cosec}^2\theta = 2$$

Thus, option (C) 2 is the correct answer.

**31. If  $f(x) = \cos^2x + \sec^2x$ , then**

**A.  $f(x) < 1$**

**B.  $f(x) = 1$**

**C.  $2 < f(x) < 1$**

**D.  $f(x) \geq 2$**

[Hint: A.M  $\geq$  G.M.]

**Solution:**

D.  $f(x) \geq 2$

Explanation:

According to the question,

We have,  $f(x) = \cos^2x + \sec^2x$

We know that, A.M  $\geq$  G.M.

$$\Rightarrow \frac{\cos^2x + \sec^2x}{2} \geq \sqrt{\cos^2x \sec^2x}$$

$$\Rightarrow \frac{\cos^2x + \sec^2x}{2} \geq \sqrt{\cos^2x \frac{1}{\cos^2x}}$$

$$\Rightarrow \frac{\cos^2x + \sec^2x}{2} \geq 1$$

$$\Rightarrow \cos^2x + \sec^2x \geq 2$$

$$\Rightarrow f(x) \geq 2$$

Thus, option (D)  $f(x) \geq 2$  is the correct answer.

**32. If  $\tan \theta = 1/2$  and  $\tan \phi = 1/3$ , then the value of  $\theta + \phi$  is**

**A.  $\pi/6$**

**B.  $\pi$**

**C. 0**

**D.  $\pi/4$**

**Solution:**

D.  $\pi/4$

Explanation:

According to the question,

**NAGEEN GROUP**  
tamso ma jyotirgamaya

$$\tan \theta = \frac{1}{2} \text{ and } \tan \phi = \frac{1}{3}$$

We know that,

$$\tan(\theta + \phi) = \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi}$$

$$= \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \times \frac{1}{3}} = \frac{\frac{5}{6}}{\frac{5}{6}} = 1$$

$$\Rightarrow \tan(\theta + \phi) = \tan \frac{\pi}{4}$$

Thus, option (D)  $\pi/4$  is the correct answer.

**33. Which of the following is not correct?**

A.  $\sin \theta = -1/5$

B.  $\cos \theta = 1$

C.  $\sec \theta = 1/2$

D.  $\tan \theta = 20$

**Solution:**

C.  $\sec \theta = 1/2$

Explanation:

According to the question,

We know that,

a)  $\sin \theta = -1/5$  is correct since  $\sin \theta \in [-1, 1]$

b)  $\cos \theta = 1$  is correct since  $\cos \theta \in [-1, 1]$

c)  $\sec \theta = 1/2$

$$\Rightarrow (1/\cos \theta) = 1/2$$

$\Rightarrow \cos \theta = 2$  is incorrect since  $\cos \theta \in [-1, 1]$

d)  $\tan \theta = 20$  is correct since  $\tan \theta \in \mathbb{R}$ .

Thus, option (C)  $\sec \theta = 1/2$  is the correct answer.

**34. The value of  $\tan 1^\circ \tan 2^\circ \tan 3^\circ \dots \tan 89^\circ$  is**

A. 0

B. 1

C.  $1/2$

D. Not defined

**Solution:**

B. 1

Explanation:

According to the question,

$$\tan 1^\circ \tan 2^\circ \tan 3^\circ \dots \tan 89^\circ$$

$$= \tan 1^\circ \tan 2^\circ \dots \tan 45^\circ \tan (90-44^\circ) \tan (90-43^\circ) \dots \tan (90-1^\circ)$$

$$= \tan 1^\circ \tan 2^\circ \dots \tan 45^\circ \cot 44^\circ \cot 43^\circ \dots \cot 1^\circ [\because \tan (90-\theta) = \cot \theta]$$

$$= \tan 1^\circ \cot 1^\circ \tan 2^\circ \cot 2^\circ \dots \tan 45^\circ \dots \tan 89^\circ \cot 89^\circ$$

$$= 1.1 \dots 1 = 1$$

Thus, option (B) 1 is the correct answer.

**35. The value of  $(1 - \tan^2 15^\circ)/(1 + \tan^2 15^\circ)$  is**

- A. 1
- B.  $\sqrt{3}$
- C.  $\sqrt{3}/2$
- D. 2

**Solution:**

C.  $\sqrt{3}/2$

Explanation:

According to the question,

Let  $\theta = 15^\circ \Rightarrow 2\theta = 30^\circ$

Now, since we know that,

$$\begin{aligned}\cos 2\theta &= \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \\ \Rightarrow \cos 30^\circ &= \frac{1 - \tan^2 15^\circ}{1 + \tan^2 15^\circ} \\ \Rightarrow \frac{\sqrt{3}}{2} &= \frac{1 - \tan^2 15^\circ}{1 + \tan^2 15^\circ}\end{aligned}$$

Thus, option (C)  $\sqrt{3}/2$  is the correct answer.

**36. The value of  $\cos 1^\circ \cos 2^\circ \cos 3^\circ \dots \cos 179^\circ$  is**

- A.  $1/\sqrt{2}$
- B. 0
- C. 1
- D. -1

**Solution:**

B. 0

Explanation:

According to the question,

Since  $\cos 90^\circ = 0$

We get,

$$\Rightarrow \cos 1^\circ \cos 2^\circ \cos 3^\circ \dots \cos 90^\circ \dots \cos 179^\circ = 0$$

Thus, option (B) 0 is the correct answer.

**37. If  $\tan \theta = 3$  and  $\theta$  lies in third quadrant, then the value of  $\sin \theta$  is**

- A.  $1/\sqrt{10}$
- B.  $-1/\sqrt{10}$
- C.  $-3/\sqrt{10}$
- D.  $3/\sqrt{10}$

**Solution:**

C.  $-3/\sqrt{10}$

Explanation:

According to the question,

Given that,  $\tan \theta = 3$  and  $\theta$  lies in third quadrant

$$\Rightarrow \cot \theta = 1/3$$



We know that,

$$\operatorname{Cosec}^2 \theta = 1 + \cot^2 \theta$$

$$= 1 + \left(\frac{1}{3}\right)^2 = 1 + \frac{1}{9} = \frac{10}{9}$$

$$\Rightarrow \sin^2 \theta = \frac{9}{10}$$

$$\Rightarrow \sin \theta = \pm \frac{3}{\sqrt{10}}$$

$$\Rightarrow \sin \theta = -\frac{3}{\sqrt{10}}, \text{ since } \theta \text{ lies in third quadrant.}$$

Thus, option (C)  $-3/\sqrt{10}$  is the correct answer.

**38. The value of  $\tan 75^\circ - \cot 75^\circ$  is equal to**

A.  $2\sqrt{3}$

B.  $2 + \sqrt{3}$

C.  $2 - \sqrt{3}$

D. 1

**Solution:**

A.  $2\sqrt{3}$

Explanation:

According to the question,

We have,

$$\tan 75^\circ - \cot 75^\circ$$

$$= \frac{\sin 75^\circ}{\cos 75^\circ} - \frac{\cos 75^\circ}{\sin 75^\circ}$$

$$= \frac{\sin^2 75^\circ - \cos^2 75^\circ}{\cos 75^\circ \sin 75^\circ}$$

$$= \frac{2(\sin^2 75^\circ - \cos^2 75^\circ)}{2 \cos 75^\circ \sin 75^\circ}$$

$$= \frac{-2 \cos 150^\circ}{\sin 150^\circ}$$

$$= -2 \cot 150^\circ$$

$$= -2 \cot (180^\circ - 30^\circ)$$

$$= 2 \cot 30^\circ$$

$$= 2\sqrt{3}$$

Thus, option (A)  $2\sqrt{3}$  is the correct answer.

# CLICK TO DOWNLOAD NCERT EXEMPLAR SOLUTIONS

**CLASS 11: MATHEMATICS (ALL CHAPTERS)**





# JOIN OUR WHATSAPP GROUPS

FOR FREE EDUCATIONAL  
RESOURCES

---





## JOIN SCHOOL OF EDUCATORS WHATSAPP GROUPS FOR FREE EDUCATIONAL RESOURCES

We are thrilled to introduce the School of Educators WhatsApp Group, a platform designed exclusively for educators to enhance your teaching & Learning experience and learning outcomes. Here are some of the key benefits you can expect from joining our group:

### BENEFITS OF SOE WHATSAPP GROUPS

---

- **Abundance of Content:** Members gain access to an extensive repository of educational materials tailored to their class level. This includes various formats such as PDFs, Word files, PowerPoint presentations, lesson plans, worksheets, practical tips, viva questions, reference books, smart content, curriculum details, syllabus, marking schemes, exam patterns, and blueprints. This rich assortment of resources enhances teaching and learning experiences.
- **Immediate Doubt Resolution:** The group facilitates quick clarification of doubts. Members can seek assistance by sending messages, and experts promptly respond to queries. This real-time interaction fosters a supportive learning environment where educators and students can exchange knowledge and address concerns effectively.
- **Access to Previous Years' Question Papers and Topper Answers:** The group provides access to previous years' question papers (PYQ) and exemplary answer scripts of toppers. This resource is invaluable for exam preparation, allowing individuals to familiarize themselves with the exam format, gain insights into scoring techniques, and enhance their performance in assessments.

- **Free and Unlimited Resources:** Members enjoy the benefit of accessing an array of educational resources without any cost restrictions. Whether its study materials, teaching aids, or assessment tools, the group offers an abundance of resources tailored to individual needs. This accessibility ensures that educators and students have ample support in their academic endeavors without financial constraints.
- **Instant Access to Educational Content:** SOE WhatsApp groups are a platform where teachers can access a wide range of educational content instantly. This includes study materials, notes, sample papers, reference materials, and relevant links shared by group members and moderators.
- **Timely Updates and Reminders:** SOE WhatsApp groups serve as a source of timely updates and reminders about important dates, exam schedules, syllabus changes, and academic events. Teachers can stay informed and well-prepared for upcoming assessments and activities.
- **Interactive Learning Environment:** Teachers can engage in discussions, ask questions, and seek clarifications within the group, creating an interactive learning environment. This fosters collaboration, peer learning, and knowledge sharing among group members, enhancing understanding and retention of concepts.
- **Access to Expert Guidance:** SOE WhatsApp groups are moderated by subject matter experts, teachers, or experienced educators can benefit from their guidance, expertise, and insights on various academic topics, exam strategies, and study techniques.

Join the School of Educators WhatsApp Group today and unlock a world of resources, support, and collaboration to take your teaching to new heights. To join, simply click on the group links provided below or send a message to +91-95208-77777 expressing your interest.

**Together, let's empower ourselves & Our Students and  
inspire the next generation of learners.**

**Best Regards,  
Team  
School of Educators**



# Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

## Kindergarten to Class XII (For Teachers Only)



[Click Here to Join](#)

**Class 1**



[Click Here to Join](#)

**Class 2**



[Click Here to Join](#)

**Class 3**



[Click Here to Join](#)

**Class 4**



[Click Here to Join](#)

**Class 5**



[Click Here to Join](#)

**Class 6**



[Click Here to Join](#)

**Class 7**



[Click Here to Join](#)

**Class 8**



[Click Here to Join](#)

**Class 9**



[Click Here to Join](#)

**Class 10**



[Click Here to Join](#)

**Class 11 (Science)**



[Click Here to Join](#)

**Class 11 (Humanities)**



[Click Here to Join](#)

**Class 11 (Commerce)**



[Click Here to Join](#)

**Class 12 (Science)**



[Click Here to Join](#)

**Class 12 (Humanities)**



[Click Here to Join](#)

**Class 12 (Commerce)**



[Click Here to Join](#)

**Kindergarten**



# Subject Wise Secondary and Senior Secondary Groups (IX & X For Teachers Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi-A



[Click Here to Join](#)

IT Code-402



[Click Here to Join](#)

Hindi-B



[Click Here to Join](#)

Artificial Intelligence

## Senior Secondary Groups (XI & XII For Teachers Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

**Geography**



[Click Here to Join](#)

**Sociology**



[Click Here to Join](#)

**Hindi Elective**



[Click Here to Join](#)

**Hindi Core**



[Click Here to Join](#)

**Home Science**



[Click Here to Join](#)

**Sanskrit**



[Click Here to Join](#)

**Psychology**



[Click Here to Join](#)

**Political Science**



[Click Here to Join](#)

**Painting**



[Click Here to Join](#)

**Vocal Music**



[Click Here to Join](#)

**Comp. Science**



[Click Here to Join](#)

**IP**



[Click Here to Join](#)

**Physical Education**



[Click Here to Join](#)

**APP. Mathematics**



[Click Here to Join](#)

**Legal Studies**



[Click Here to Join](#)

**Entrepreneurship**



[Click Here to Join](#)

**French**



[Click Here to Join](#)

**IT**



[Click Here to Join](#)

**Artificial Intelligence**

## **Other Important Groups (For Teachers & Principal's)**



[Click Here to Join](#)

**Principal's Group**



[Click Here to Join](#)

**Teachers Jobs**



[Click Here to Join](#)

**IIT/NEET**

# Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

## Kindergarten to Class XII (For Students Only)



[Click Here to Join](#)

**Class 1**



[Click Here to Join](#)

**Class 2**



[Click Here to Join](#)

**Class 3**



[Click Here to Join](#)

**Class 4**



[Click Here to Join](#)

**Class 5**



[Click Here to Join](#)

**Class 6**



[Click Here to Join](#)

**Class 7**



[Click Here to Join](#)

**Class 8**



[Click Here to Join](#)

**Class 9**



[Click Here to Join](#)

**Class 10**



[Click Here to Join](#)

**Class 11 (Science)**



[Click Here to Join](#)

**Class 11 (Humanities)**



[Click Here to Join](#)

**Class 11 (Commerce)**



[Click Here to Join](#)

**Class 12 (Science)**



[Click Here to Join](#)

**Class 12 (Humanities)**



[Click Here to Join](#)

**Class 12 (Commerce)**



[Click Here to Join](#)

**Artificial Intelligence  
(VI TO VIII)**

# Subject Wise Secondary and Senior Secondary Groups (IX & X For Students Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi



[Click Here to Join](#)

IT Code



[Click Here to Join](#)

Artificial Intelligence

## Senior Secondary Groups (XI & XII For Students Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

**Geography**



[Click Here to Join](#)

**Sociology**



[Click Here to Join](#)

**Hindi Elective**



[Click Here to Join](#)

**Hindi Core**



[Click Here to Join](#)

**Home Science**



[Click Here to Join](#)

**Sanskrit**



[Click Here to Join](#)

**Psychology**



[Click Here to Join](#)

**Political Science**



[Click Here to Join](#)

**Painting**



[Click Here to Join](#)

**Music**



[Click Here to Join](#)

**Comp. Science**



[Click Here to Join](#)

**IP**



[Click Here to Join](#)

**Physical Education**



[Click Here to Join](#)

**APP. Mathematics**



[Click Here to Join](#)

**Legal Studies**



[Click Here to Join](#)

**Entrepreneurship**



[Click Here to Join](#)

**French**



[Click Here to Join](#)

**IT**



[Click Here to Join](#)

**AI**



[Click Here to Join](#)

**IIT/NEET**



[Click Here to Join](#)

**CUET**

## Groups Rules & Regulations:

**To maximize the benefits of these WhatsApp groups, follow these guidelines:**

1. Share your valuable resources with the group.
2. Help your fellow educators by answering their queries.
3. Watch and engage with shared videos in the group.
4. Distribute WhatsApp group resources among your students.
5. Encourage your colleagues to join these groups.

### **Additional notes:**

1. Avoid posting messages between 9 PM and 7 AM.
2. After sharing resources with students, consider deleting outdated data if necessary.
3. It's a NO Nuisance groups, single nuisance and you will be removed.
  - No introductions.
  - No greetings or wish messages.
  - No personal chats or messages.
  - No spam. Or voice calls
  - Share and seek learning resources only.

**Please only share and request learning resources. For assistance, contact the helpline via WhatsApp: +91-95208-77777.**



# Join Premium WhatsApp Groups Ultimate Educational Resources!!

Join our premium groups and just Rs. 1000 and gain access to all our exclusive materials for the entire academic year. Whether you're a student in Class IX, X, XI, or XII, or a teacher for these grades, Artham Resources provides the ultimate tools to enhance learning. Pay now to delve into a world of premium educational content!

[Click here for more details](#)



**Click Here to Join**

**Class 9**



**Click Here to Join**

**Class 10**



**Click Here to Join**

**Class 11**



**Click Here to Join**

**Class 12**

📢 Don't Miss Out! Elevate your academic journey with top-notch study materials and secure your path to top scores! Revolutionize your study routine and reach your academic goals with our comprehensive resources. Join now and set yourself up for success! 🇧🇩🌟

**Best Wishes,**

**Team**

**School of Educators & Artham Resources**

# SKILL MODULES BEING OFFERED IN MIDDLE SCHOOL



Artificial Intelligence



Beauty & Wellness




Design Thinking & Innovation



Financial Literacy



Handicrafts



Information Technology



Marketing/Commercial Application




Mass Media - Being Media Literate




Travel & Tourism



Coding




Data Science (Class VIII only)




Augmented Reality / Virtual Reality



Digital Citizenship




Life Cycle of Medicine & Vaccine



Things you should know about keeping Medicines at home




What to do when Doctor is not around



Humanity & Covid-19




Blue Pottery




Pottery



Block Printing




Food



Food Preservation




Baking



Herbal Heritage



Khadi




Mask Making



Mass Media



Making of a Graphic Novel




Kashmiri Embroidery



Embroidery




Rockets



Satellites




Application of Satellites




Photography

# SKILL SUBJECTS AT SECONDARY LEVEL (CLASSES IX – X)



Retail



Information Technology




Security




Automotive



Introduction To Financial Markets



Introduction To Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking & Insurance



Marketing & Sales



Health Care



Apparel



Multi Media



Multi Skill Foundation Course



Artificial Intelligence



Physical Activity Trainer



Data Science



Electronics & Hardware (NEW)



Foundation Skills For Sciences (Pharmaceutical & Biotechnology)(NEW)



Design Thinking & Innovation (NEW)



# SKILL SUBJECTS AT SR. SEC. LEVEL (CLASSES XI – XII)



Retail



Information Technology



Web Application



Automotive



Financial Markets Management



Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking



Marketing




Health Care



Insurance




Horticulture



Typography & Comp.  
Application



Geospatial Technology



Electrical Technology



Electronic Technology



Multi-Media




Taxation



Cost Accounting



Office Procedures & Practices



Shorthand (English)




Shorthand (Hindi)



Air-Conditioning & Refrigeration



Medical Diagnostics



Textile Design




Design



Salesmanship



Business Administration



Food Nutrition & Dietetics



Mass Media Studies



Library & Information Science



Fashion Studies



Applied Mathematics



Yoga



Early Childhood Care & Education




Artificial Intelligence



Data Science



Physical Activity Trainer(new)



Land Transportation Associate (NEW)



Electronics & Hardware (NEW)



Design Thinking & Innovation (NEW)